

February 1, 2008

GROWTH LAWS FOR PHASE ORDERING

A. J. Bray and A. D. Rutenberg

Department of Theoretical Physics

The University, Manchester M13 9PL

bray@mphv2.phy.man.ac.uk, ar@mphv2.phy.man.ac.uk

ABSTRACT

We determine the characteristic length scale, $L(t)$, in phase ordering kinetics for both scalar and vector fields, with either short- or long-range interactions, and with or without conservation laws. We obtain $L(t)$ consistently by comparing the global rate of energy change to the energy dissipation from the local evolution of the order parameter. We derive growth laws for $O(n)$ models, and our results can be applied to other systems with similar defect structures.

cond-mat/9303011

Systems quenched from a disordered phase into an ordered phase do not order instantaneously. Instead, the length scale of ordered regions grows at a characteristic rate as different broken symmetry phases compete to select the ground state. During this process, energy is dissipated and topological defects, if present, are destroyed. Traditionally, systems with scalar order parameters, such as binary alloys and Ising models, have been studied [1, 2]. In such systems domains of both phases grow and the intervening domain walls, which are the characteristic topological defects of scalar systems, shrink in total area; the energy of the system is dissipated as the total area of domain wall decreases. Recently, there has been a growing interest in systems with vector and more complex order parameters which have defect structures such as lines, points, and textures. An exciting series of experiments [3, 4] and simulations [5, 6, 7, 8, 9] has been performed to explore these systems. Of particular interest is the time dependence, $L(t)$, of the characteristic length scale. This growth law provides a basis for further analysis and a fundamental test of approximation schemes. Growth laws have been obtained in individual cases [10, 11, 12, 13], but the variety of possible systems has prevented the development of a unified approach. Renormalization group (RG) ideas [12, 14] have been used to predict the leading power-law dependence of growth laws, but are not easy to apply in systems without conservation laws. An approach to calculating growth laws, including logarithmic factors, for a wide variety of systems would be welcome.

In this Letter we determine the characteristic length scale, $L(t)$, for quenched systems with both scalar and vector fields, with either short- or long-range interactions, and with or without conservation laws. We base our approach on the energy dissipation that occurs as the system relaxes towards its ground state. We evaluate the energy dissipation by considering topological defects, which dominate the dynamics when they exist. We compare the global rate of energy change to the

energy dissipation from the local evolution of the order parameter. For systems with a single characteristic scale describing correlations as well as dissipation we self-consistently determine the growth law, $L(t)$.

Our results are summarized in tables 1 and 2. We find that conservation laws (μ) and long-range interactions (σ) are irrelevant to the growth law below and above, respectively, values determined by the number of components (n) in the order parameter of the system. At the marginal values, logarithmic factors are introduced. The growth laws obtained are independent of the spatial dimension of the system. Our results should be compared to growth laws obtained by other means. Heuristic arguments yield growth laws of $L \sim t^{1/2}$ for non-conserved [11] and $L \sim t^{1/3}$ for conserved [10] scalar order parameters. RG arguments [12] reproduce the $t^{1/3}$ law for conserved scalar fields, predict $L \sim t^{1/4}$ for a conserved vector field, and also treat conserved systems with long-range interactions [14]. Simulations [5] suggest a $L \sim t^{1/2}$ growth law for non-conserved vector fields, and simulation [6] and experiments [3] obtain the same law for the related tensor fields that describe nematic liquid crystals. This result has also been obtained by approximate treatments for general n [15] and verified in the large- n limit [13]. Our approach provides a unified theoretical basis for the leading power-law behavior suggested by these results, and also determines any logarithmic factors.

The dynamic scaling hypothesis describes most systems in the late stages of growth [16]. Accordingly, the correlation function of the order parameter, $C(\mathbf{r}, t) = \langle \vec{\phi}(\mathbf{x}, t) \cdot \vec{\phi}(\mathbf{x} + \mathbf{r}, t) \rangle$, with an average over initial conditions, should exhibit the scaling form $C(\mathbf{r}, t) = f(r/L(t))$, with a single characteristic length scale $L(t)$. Fourier transforming this, we obtain the scaling form for the structure factor $S(\mathbf{k}, t) = \langle \vec{\phi}_{\mathbf{k}}(t) \cdot \vec{\phi}_{-\mathbf{k}}(t) \rangle$,

$$S(\mathbf{k}, t) = L(t)^d g(kL(t)) , \quad (1)$$

where d is the spatial dimension. A generic energy functional of such a system, with an n -component field, $\vec{\phi}(\vec{x})$, is

$$H[\vec{\phi}] = \int d^d x [(\nabla \vec{\phi})^2 + V(\vec{\phi})] , \quad (2)$$

where $V(\vec{\phi})$ is a “mexican hat” shaped potential such as $V(\vec{\phi}) = (\vec{\phi}^2 - 1)^2$. The equation of motion for the ordering kinetics of the Fourier components $\vec{\phi}_{\mathbf{k}}$ [17] is

$$\partial_t \vec{\phi}_{\mathbf{k}} = -k^\mu (\partial H / \partial \vec{\phi}_{-\mathbf{k}}), \quad (3)$$

where we work at temperature $T = 0$, with no thermal noise, since T appears to be an ‘irrelevant variable’ for ordering kinetics within the ordered phase [12, 13]. The conventional non-conserved (model A dynamics) and conserved (model B dynamics) cases are $\mu = 0$ and $\mu = 2$, respectively.

Integrating the rate of energy dissipation from each Fourier mode, and then using the equation of motion (3), we find

$$\begin{aligned} d\epsilon/dt &= \int_{\mathbf{k}} \langle (\partial H / \partial \vec{\phi}_{\mathbf{k}}) \cdot \partial_t \vec{\phi}_{\mathbf{k}} \rangle \\ &= - \int_{\mathbf{k}} k^{-\mu} \langle \partial_t \vec{\phi}_{\mathbf{k}} \cdot \partial_t \vec{\phi}_{-\mathbf{k}} \rangle , \end{aligned} \quad (4)$$

where $\epsilon = \langle H \rangle / V$ is the mean energy density, and $\int_{\mathbf{k}}$ is the momentum integral $\int d^d k / (2\pi)^d$. We will relate the scaling behavior of both sides of (4) to that of appropriate integrals over the structure factor, $S(\mathbf{k}, t)$. Either the integrals converge and the dependence on the scale $L(t)$ can be extracted using the scaling form (1), or a divergent contribution dominates the integral. Any divergence will occur because of small-scale structure at $kL \gg 1$, since no structure has formed yet for $kL \ll 1$. This small-scale structure will only come from topological defects, so we expect $S(\mathbf{k}, t)$ to be proportional to the density of defect core, $\rho_{\text{def}} \sim L^{d-n}/L^d \sim L^{-n}$ for $kL \gg 1$ [18]. The scaling form (1) then implies

$$S(\mathbf{k}, t) \sim L^{-n} k^{-(d+n)} , \quad kL \gg 1, \quad (5)$$

which is a generalized Porod's law for $n \leq d$ [19, 20].

We first calculate the scaling behavior of the energy density, ϵ , which is captured by that of the gradient term in (2):

$$\begin{aligned}\epsilon &\sim \langle (\nabla \vec{\phi})^2 \rangle \\ &= \int_{\mathbf{k}} k^2 L^d g(kL) ,\end{aligned}\tag{6}$$

where we have used the scaling form (1) for the structure factor. If the integral converges, then a simple change of variables yields $\epsilon \sim L^{-2}$. If the integral diverges for $kL \gg 1$, then we use Porod's law (5) and impose a cutoff at $k \sim 1/\xi$, where ξ is the defect core size. We obtain [20]

$$\begin{aligned}\epsilon &\sim L^{-n} \xi^{n-2} , & n < 2 , \\ &\sim L^{-2} \ln(L/\xi) , & n = 2 , \\ &\sim L^{-2} , & n > 2 .\end{aligned}\tag{7}$$

We see that the energy is dominated by the defect core density ρ_{def} for $n < 2$, by the defect field at all length scales for $n = 2$, and by variations of the order parameter at scale $L(t)$ for $n > 2$.

We now evaluate the right side of (4) in a similar way. Using the scaling assumption for the two-time function, $\langle \vec{\phi}_{\mathbf{k}}(t) \cdot \vec{\phi}_{-\mathbf{k}}(t') \rangle = k^{-d} g(kL(t), kL(t'))$, we find

$$\begin{aligned}\langle \partial_t \vec{\phi}_{\mathbf{k}} \cdot \partial_t \vec{\phi}_{-\mathbf{k}} \rangle &= \left. \frac{\partial^2}{\partial t \partial t'} \right|_{t=t'} \langle \vec{\phi}_{\mathbf{k}}(t) \cdot \vec{\phi}_{-\mathbf{k}}(t') \rangle \\ &= (dL/dt)^2 L^{d-2} h(kL).\end{aligned}\tag{8}$$

When the momentum integral on the right of (4) converges, we obtain, using (8), $d\epsilon/dt \sim -(dL/dt)^2 L^{\mu-2}$. However, if the integral diverges we need to know the behaviour of (8) for $kL \gg 1$.

In this regime, where the important structure is short compared to the typical interdefect spacing, the scaling functions are determined by the field of a single

defect [18], and the defect field will be comoving with the defect core,

$$\partial_t \vec{\phi} = \vec{\omega} \times \vec{\phi} - \mathbf{v} \cdot \nabla \vec{\phi} . \quad (9)$$

Here, $\mathbf{v}$ is the velocity of the defect and $\vec{\omega}$ represents rotations for vector fields. For $kL \gg 1$, the velocity term dominates because the rotation scales as $\omega \sim (dL/dt)/L$ while the gradient term involves the short-scale structure, $\mathbf{v} \cdot \nabla \sim (dL/dt)k$. In estimating $v \sim (dL/dt)$, we are assuming that dissipation occurs by the evolution of defects of scale $L(t)$. Using $\partial_t \vec{\phi}_{\mathbf{k}} \sim (dL/dt)k \vec{\phi}_{\mathbf{k}}$ and Porod's law (5), we have

$$\langle \partial_t \vec{\phi}_{\mathbf{k}} \cdot \partial_t \vec{\phi}_{-\mathbf{k}} \rangle \sim L^{-n} k^{-(d+n-2)} (dL/dt)^2 , \quad kL \gg 1 . \quad (10)$$

Using (10) in the right of (4) shows that the integral is convergent for $kL \gg 1$ when $n + \mu > 2$. Otherwise the integral is dominated by k near the upper cut-off $1/\xi$. We obtain

$$\begin{aligned} \int_{\mathbf{k}} k^{-\mu} \langle \partial_t \vec{\phi}_{\mathbf{k}} \cdot \partial_t \vec{\phi}_{-\mathbf{k}} \rangle &\sim L^{-n} \xi^{n+\mu-2} (dL/dt)^2 , \quad n + \mu < 2 , \\ &\sim L^{-n} \ln(L/\xi) (dL/dt)^2 , \quad n + \mu = 2 , \\ &\sim L^{\mu-2} (dL/dt)^2 , \quad n + \mu > 2 . \end{aligned} \quad (11)$$

In what cases does dissipation occur primarily at scales $L(t)$, as we have just assumed? For $n > 2$, the energy density (7), and hence dissipation, is dominated by variations at scale $L(t)$. For $n \leq 2$, with $d > n$, the energy density is proportional to the defect core volume, but the dissipation is still dominated by scales of order $L(t)$. For this case the rate of energy dissipation at a length scale l is given by the rate at which the core volume disappears at that scale. So the total energy dissipation is given by

$$d\epsilon/dt \sim \int dl n(l) \dot{l} l^{d-n-1} , \quad (12)$$

where $\dot{l}$ is the rate of evolution of defect structures of scale l , $n(l)$ is the number density of features of scale l , and $\dot{l} l^{d-n-1}$ is the rate of change of the defect core

volume at scale l . But $j(l) \sim n(l) \dot{l}$ is just the number flux of defect features, and imposing the continuity equation we see that $\dot{N} \sim j(0)$, where N is the average density of defects. N scales as a volume, $N \sim 1/L^d$, and so $\dot{N}$ is a constant for times of order $L/\dot{L}$. This implies that $j(l)$ is constant for $l \ll L$, in order to provide a constant rate of defect extinction. We see that $d\epsilon/dt \sim \int dl l^{d-n-1}$ for $l \ll L$, and the convergence at small l for $d > n$ implies that structures with scales $l \sim L(t)$ dominate the energy dissipation.

For $n = d$ with $n < 2$, the marginal case in the above argument, the energy density (7) is dominated by the core energy of point defects and dissipation is dominated by defect pairs annihilating. Since the dissipation occurs at separations $l \sim \xi \ll L$ we do not expect our approach to cover these cases. In fact, since the bulk energy density does not depend on defect separations $l \gg \xi$, we expect the system to be disordered, with an equilibrium density of defects, at any non-zero temperature. At $T = 0$ we expect slow growth laws that depend on the details of the potential $V(\vec{\phi})$. These cases, $n = d$ with $n < 2$, are at their lower critical dimension.

The $2d$ XY model is outside the scope of this Letter, since a more detailed argument [21] shows that dissipation occurs significantly at all length scales between ξ and $L(t)$. Naively, we might expect scaling violations for this case. Yurke *et al.* [9] suggest a growth law of $L \sim (t/\ln t)^{1/2}$ for non-conserved order-parameters, while simulations by Mondello and Goldenfeld [7] find $L \sim t^{1/4}$ for the conserved case.

In the cases where dissipation occurs at the scale of $L(t)$, we compare the rate of energy dissipation between (11) and the time derivative of (7) to obtain dL/dt and hence $L(t)$. The results are summarized in table 1.

For non-conserved fields ($\mu = 0$), we find $L \sim t^{1/2}$ for all systems (with $d > n$ or $n > 2$). Leading corrections in the $n = 2$ case are interesting: the $\ln L$ factors in (13) and (17) will in general have different effective cutoffs, of order the core size ξ .

This leads to a logarithmic correction to scaling, $L \sim t^{1/2}(1 + O(1/\ln t))$, and may account for the smaller exponent (~ 0.45) seen in simulations of $O(2)$ systems [5, 6].

For conserved fields ($\mu > 0$) our results agree with the the RG analysis [12], with additional logarithmic factors for the marginal cases $n = 2$ and $n + \mu = 2$. We see that the conservation law is only relevant for $n + \mu \geq 2$. Therefore for vector fields ($n \geq 2$), any $\mu > 0$ is sufficient to change the growth law, while for scalar fields ($n = 1$) the conservation law is irrelevant for $\mu < 1$, in agreement with the RG analysis [12]. Simulations by Siegert and Rao [8] for $n = 2$, with $d = 3$ and $\mu = 2$, obtain growth exponents slightly over $1/4$, which may be difficult to distinguish from our predicted $L \sim (t \ln t)^{1/4}$ behavior.

The above discussion covers systems with purely short-ranged interactions. We can also add long-ranged interactions to our energy functional (2) [22, 14]:

$$H_{LR} \sim \int d^d x \int d^d r \frac{(\vec{\phi}(\mathbf{x} + \mathbf{r}) - \vec{\phi}(\mathbf{x}))^2}{r^{d+\sigma}}. \quad (13)$$

The evaluation of the right side of (4) is unchanged, but the mean energy is now given by

$$\epsilon_{LR} \sim \int_{\mathbf{k}} k^\sigma S(\mathbf{k}, t), \quad \sigma < 2. \quad (14)$$

We consider $\sigma < 2$ only, since for $\sigma > 2$ the energy is dominated by short distances and we recover the short-range k^2 form [23]. Inserting the scaling form (1) for $S(\mathbf{k}, t)$, and noting the asymptotic behavior (5), gives

$$\begin{aligned} \epsilon_{LR} &\sim L^{-n} \xi^{n-\sigma}, & n < \sigma, \\ &\sim L^{-\sigma} \ln(L/\xi), & n = \sigma, \\ &\sim L^{-\sigma} & n > \sigma. \end{aligned} \quad (15)$$

We then compare the rate of energy dissipation between (11), which is unchanged, and the time derivative of (15), to find dL/dt and generate table 2. Our results agree

with the leading power laws from an RG treatment by Bray [14] for the conserved case ($\mu > 0$), while here we obtain the growth laws for all cases, with any logarithmic corrections.

From (15), we see that the long-range interactions are relevant only for $n \geq \sigma$ [14]. This correctly indicates that defects are non-interacting for $n < \sigma$, where the dynamics are driven by a generalized surface tension. Note that cases with $\sigma < 0$ lead to an instability at fixed non-zero wave-vector which breaks the scaling (1) and hence our approach does not directly apply. It is also interesting to note that naive power counting using H_{LR} (13) in the dynamics (3) yields the incorrect growth law $L \sim t^{1/\sigma}$ for non-conserved scalar fields. The fact that power counting obtains the correct growth law, $L \sim t^{1/2}$, for short range interactions with $n \leq 2$, can be seen as a fortuitous balancing of the UV divergences in (6) and (8).

Since systems without topological defects ($n > d + 1$) will have convergent momentum integrals for $kL \gg 1$, we obtain $L \sim t^{1/(\sigma+\mu)}$, where $\sigma = 2$ for short-ranged forces. We can also apply this result to systems with topological textures ($n = d+1$), even though the appropriate Porod's law is not known. Since defects with $n > d$ must be spatially extended and without a core, they will have a smaller large- k tail to their structure factor $S(\mathbf{k}, t)$ than any defects with cores. So for $n > 2$, when the energy dissipation clearly occurs at length scales of order $L(t)$ (see (7)) and the momentum integrals for defects with cores converge, our results should apply. As a result, tables 1 and 2 will apply for *any* system, except perhaps those with $d \leq n \leq 2$ [24]. Of course, whether the structure factor scales, as in (1), for systems with textures is an open question.

The strength of our approach is that it can be applied to systems with more complicated order parameters than n -component vectors. The details of the energy functional (2) are unimportant to this approach [25]. All we need is the existence of

some short or long-ranged “elastic energy” (σ), an effective conservation law (μ), and the defect structure if any. Our derivation is independent of the initial conditions, and so, e.g., applies equally to critical and off-critical quenches *as long as the system scales at late times*. We simply choose a Porod’s law (5) to represent the dominant defect type, which is the one represented by the smallest n . For example, in bulk nematic liquid crystals, the existence of string defects leads to (5) with $n = 2$, which with no conservation law implies a $L \sim t^{1/2}$ growth law, consistent with recent experiments [3] and simulations [6].

In summary, we obtain growth laws for phase ordering based only on the dynamic scaling hypothesis and a generalized Porod’s law for the large-momentum tail of the structure factor. By focusing on the total energy dissipation, rather than the details of the system, we do not need to consider defect dynamics. This results in a powerful yet simple method that addresses many cases of interest.

We thank R. E. Blundell, D. A. Huse, H. H. Lee, and M. A. Moore for discussions.

References

- [1] For reviews see, e.g., J. D. Gunton, M. San Miguel and P. S. Sahni, in *Phase Transitions and Critical Phenomena*, Vol. 8, eds. C. Domb and J. L. Lebowitz (Academic, New York, 1983) p.267; J. S. Langer, in *Solids Far From Equilibrium*, ed. C. Godrèche (Cambridge, Cambridge, 1992).
- [2] N. Mason, A. N. Pargellis, and B. Yurke, Phys. Rev. Lett. **70**, 190 (1993).
These authors study a twisted nematic liquid crystal.
- [3] A. P. Y. Wong, P. Wiltzius and B. Yurke, Phys. Rev. Lett. **68**, 3583 (1992);
A. P. Y. Wong, P. Wiltzius, R. G. Larson and B. Yurke, preprint.

- [4] A. N. Pargellis, P. Finn, J. W. Goodby, P. Panizza, B. Yurke, and P. E. Cladis, Phys. Rev. A **46**, 7765 (1992).
- [5] For $n = 2$: H. Toyoki, J. Phys. Soc. Jpn. **60**, 1433 (1991); M. Mondello and N. Goldenfeld, Phys. Rev. A **45**, 657 (1992); for $n = 3$: H. Toyoki, J. Phys. Soc. Jpn. **60**, 1153 (1991).
- [6] R. E. Blundell and A. J. Bray, Phys. Rev. A **46**, R6154 (1992). Our results indicate that nematic liquid crystals, with line defects, should have the same growth law and leading correction as the $O(2)$ model.
- [7] M. Mondello and N. Goldenfeld, preprint.
- [8] M. Siegert and M. Rao, preprint.
- [9] B. Yurke, A. N. Pargellis, T. Kovacs and D. A. Huse, preprint. See also [4].
- [10] I. M. Lifshitz and V. V. Slyozov, J. Phys. Chem. Solids **19**, 35 (1961); D. A. Huse, Phys. Rev. B **34**, 7845 (1986).
- [11] I. M. Lifshitz, Zh. Eksp. Teor. Fiz. **42**, 1354 (1962) [Sov. Phys. – JETP **15**, 939 (1962)]; S. M. Allen and J. W. Cahn, Acta. Metall. **27**, 1085 (1979).
- [12] A. J. Bray, Phys. Rev. Lett. **62**, 2841 (1989); A. J. Bray, Phys. Rev. B **41**, 6724 (1990).
- [13] T. J. Newman and A. J. Bray, J. Phys. A **23**, 4491 (1990) and references therein.
- [14] A. J. Bray, preprint.

- [15] F. Liu and G. F. Mazenko, Phys. Rev. B **45**, 6989 (1992); A. J. Bray and K. Humayun, J. Phys. A **25**, 2191 (1992); H. Toyoki and K. Honda, Prog. Theor. Phys. **78**, 237 (1987).
- [16] H. Furukawa, Adv. Phys. **34**, 703 (1985). See also A. Coniglio and M. Zannetti, Europhys. Lett. **10**, 575 (1989) and A. J. Bray and K. Humayan, Phys. Rev. Lett. **68**, 1559 (1992).
- [17] P. C. Hohenberg and B. I. Halperin, Rev. Mod. Phys. **49**, 435 (1977).
- [18] A. J. Bray, Phys. Rev. E **47**, 228 (1993); A. J. Bray and K. Humayun, Phys. Rev. E **47**, R9, (1993).
- [19] A. J. Bray and S. Puri, Phys. Rev. Lett. **67**, 2670 (1991).
- [20] H. Toyoki, Phys. Rev. B **45**, 1965 (1992).
- [21] A. J. Bray and A. D. Rutenberg, unpublished.
- [22] H. Hayakawa, Z. Rácz and T. Tsuzuki, preprint; A. J. Bray, J. Phys. C **19**, 6225 (1986).
- [23] The short-range growth laws correspond to the limit $\sigma \rightarrow 2$.
- [24] T. J. Newman, A. J. Bray, and M. A. Moore, Phys. Rev. B **42** 4514, (1990) found $L \sim t^{1/4}$ for $d = 1$, $n = 2$, and $\mu = 0$.
- [25] Of course this means that our approach will not address systems with a potential-dependent growth law, e.g. $d = n$ for $n < 2$. We also do not address quenches in which thermal noise is essential, such as systems with static disorder (see D. A. Huse and C. L. Henley, Phys. Rev. Lett. **54**, 2708 (1985)), or quenches to a $T > 0$ critical point.

$L(t)$	$n < 2$	$n = 2$	$n > 2$
$n + \mu < 2$	$t^{1/2}$	—	—
$n + \mu = 2$	$(t/\ln t)^{1/2}$	$t^{1/2}$	—
$n + \mu > 2$	$t^{1/(n+\mu)}$	$(t \ln t)^{1/(2+\mu)}$	$t^{1/(2+\mu)}$

Table 1: The growth law of the length scale $L(t)$ for various number of components, n , and conservation laws, μ . Note that $n = d$ is excluded for $n \leq 2$.

$L(t)$	$n < \sigma$	$n = \sigma$	$n > \sigma$
$n + \mu < 2$	$t^{1/2}$	$(t \ln t)^{1/2}$	$t^{1/(2+\sigma-n)}$
$n + \mu = 2$	$(t/\ln t)^{1/2}$	$t^{1/2}$	$(t/\ln t)^{1/(2+\sigma-n)}$
$n + \mu > 2$	$t^{1/(n+\mu)}$	$(t \ln t)^{1/(\sigma+\mu)}$	$t^{1/(\sigma+\mu)}$

Table 2: The growth law of the length scale $L(t)$ for long-range forces, $0 < \sigma < 2$, with various number of components, n , and conservation laws, μ . Note that $n = d$ is excluded for $n \leq \sigma$.